

COVID-19 Information[Public health information \(CDC\)](#)[Research information \(NIH\)](#)[SARS-CoV-2 data \(NCBI\)](#)[Prevention and treatment information \(HHS\)](#)[Español](#)

FULL TEXT LINKS

Review [Lasers Med Sci.](#) 2015 Feb;30(2):925-39. doi: 10.1007/s10103-013-1465-4.

Epub 2013 Nov 19.

Effect of phototherapy (low-level laser therapy and light-emitting diode therapy) on exercise performance and markers of exercise recovery: a systematic review with meta-analysis

[Ernesto Cesar Pinto Leal-Junior](#)¹, [Adriane Aver Vanin](#), [Eduardo Foschini Miranda](#), [Paulo de Tarso Camillo de Carvalho](#), [Simone Dal Corso](#), [Jan Magnus Bjordal](#)

Affiliations

PMID: 24249354 DOI: [10.1007/s10103-013-1465-4](#)

Abstract

Recent studies have explored if phototherapy with low-level laser therapy (LLLT) or narrow-band light-emitting diode therapy (LEDT) can modulate activity-induced skeletal muscle fatigue or subsequently protect against muscle injury. We performed a systematic review with meta-analysis to investigate the effects of phototherapy applied before, during and after exercises. A literature search was performed in Pubmed/Medline database for randomized controlled trials (RCTs) published from 2000 through 2012. Trial quality was assessed with the ten-item PEDro scale. Main outcome measures were selected as: number of repetitions and time until exhaustion for muscle performance, and creatine kinase (CK) activity to evaluate risk for exercise-induced muscle damage. The literature search resulted in 16 RCTs, and three articles were excluded due to poor quality assessment scores. From 13 RCTs with acceptable methodological quality (≥ 6 of 10 items), 12 RCTs irradiated phototherapy before exercise, and 10 RCTs reported significant improvement for the main outcome measures related to performance. The time until exhaustion increased significantly compared to placebo by 4.12 s (95% CI 1.21-7.02, $p < 0.005$) and the number of repetitions increased by 5.47 (95% CI 2.35-8.59, $p < 0.0006$) after phototherapy. Heterogeneity in trial design and results precluded meta-analyses for biochemical markers, but a quantitative analysis showed positive results in 13 out of 16 comparisons. The most significant and consistent results were found with red or infrared wavelengths and phototherapy application before exercises, power outputs between 50 and 200 mW and doses of 5 and 6 J per point (spot). We conclude that phototherapy (with lasers and LEDs) improves muscular performance and accelerate recovery mainly when applied before exercise.

Related information

[MedGen](#)

[PubChem Compound \(MeSH Keyword\)](#)

LinkOut - more resources

Full Text Sources

[Springer](#)

Other Literature Sources

[The Lens - Patent Citations](#)

[scite Smart Citations](#)

Medical

[ClinicalTrials.gov](#)

[MedlinePlus Health Information](#)

Research Materials

[NCI CPTC Antibody Characterization Program](#)